


SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average

Condition	Category	Measure Name
Heart Attack Patients Given Aspirin at Arrival	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Heart Attack Patients Given Fibrinolytic Medication Within 30 Minutes Of Arrival	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Heart Attack Patients Given PCI Within 90 Minutes Of Arrival	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Heart Attack Patients Given ACE Inhibitor or ARB for Left Ventricular Systolic Dysfunction (LVSD)	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Heart Attack Patients Given Smoking Cessation Advice/Counseling	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Heart Attack Patients Given Aspirin at Discharge	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Heart Attack Patients Given Beta Blocker at Discharge	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Heart Attack Patients Given a Prescription for a Statin at Discharge	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Heart Failure Patients Given Discharge Instructions	Heart Failure	National Average of Hospitals submitting data:

SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average



SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average

Heart Failure Patients Given an Evaluation of Left Ventricular Systolic (LVS) Function	Heart Failure	National Average of Hospitals submitting data:
Heart Failure Patients Given ACE Inhibitor or ARB for Left Ventricular Systolic Dysfunction (LVSD)	Heart Failure	National Average of Hospitals submitting data:
Heart Failure Patients Given Smoking Cessation Advice/Counseling	Heart Failure	National Average of Hospitals submitting data:
Pneumonia Patients Given Initial Antibiotic(s) within 6 Hours After Arrival	Pneumonia	National Average of Hospitals submitting data:
Pneumonia Patients Whose Initial Emergency Room Blood Culture Was Performed Prior To The Administration Of The First Hospital Dose Of Antibiotics	Pneumonia	National Average of Hospitals submitting data:
Pneumonia Patients Given the Most Appropriate Initial Antibiotic(s)	Pneumonia	National Average of Hospitals submitting data:
Pneumonia Patients Given Smoking Cessation Advice/Counseling	Pneumonia	National Average of Hospitals submitting data:
Pneumonia Patients Assessed	Pneumonia	National Average of Hospitals

SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average

98

96

99

96

97

94

98

95

SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average

and Given Pneumococcal
Vaccination

Pneumonia Patients Assessed
and Given Influenza
Vaccination

submitting data:

National Average of Hospitals
submitting data:

Surgery patients who were
given an antibiotic at the right
time (within one hour before
surgery) to help prevent
infection

Surgical Care Improvement
Project

National Average of Hospitals
submitting data:

Surgery patients whose
preventive antibiotics were
stopped at the right time (within
24 hours after surgery)

Surgical Care Improvement
Project

National Average of Hospitals
submitting data:

Patients who got treatment at
the right time (within 24 hours
before or after their surgery) to
help prevent blood clots after
certain types of surgery

Surgical Care Improvement
Project

National Average of Hospitals
submitting data:

Surgery patients who were
taking heart drugs called beta
blockers before coming to the
hospital, who were kept on the
beta blockers during the period
just before and after their
surgery

Surgical Care Improvement
Project

National Average of Hospitals
submitting data:

SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average

93

98

97

96

96

SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average

Surgery patients who were given the right kind of antibiotic to help prevent infection	Surgical Care Improvement Project	National Average of Hospitals submitting data:
Heart surgery patients whose blood sugar (blood glucose) is kept under good control in the days right after surgery	Surgical Care Improvement Project	National Average of Hospitals submitting data:
Surgery patients needing hair removed from the surgical area before surgery, who had hair removed using a safer method (electric clippers or hair removal cream – not a razor)	Surgical Care Improvement Project	National Average of Hospitals submitting data:
Patients having surgery who were actively warmed in the operating room or whose body temperature was near normal by the end of surgery.	Surgical Care Improvement Project	National Average of Hospitals submitting data:
Surgery patients whose doctors ordered treatments to prevent blood clots after certain types of surgeries	Surgical Care Improvement Project	National Average of Hospitals submitting data:
Children Who Received Reliever Medication While Hospitalized for Asthma	Children's Asthma	National Average of Hospitals submitting data:
Children Who Received	Children's Asthma	National Average of Hospitals

SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average

98

95

100

99

97

100

100

SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average

Systemic Corticosteroid Medication (oral and IV Medication That Reduces Inflammation and Controls Symptoms) While Hospitalized for Asthma		submitting data:
Children and their Caregivers Who Received a Home Management Plan of Care Document While Hospitalized for Asthma	Children's Asthma	National Average of Hospitals submitting data:
Average number of minutes before outpatients with chest pain or possible heart attack who needed specialized care were transferred to another hospital A lower number of minutes is better	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Average number of minutes before outpatients with chest pain or possible heart attack got an ECG A lower number of minutes is better	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Outpatients with chest pain or possible heart attack who got drugs to break up blood clots	Heart Attack or Chest Pain	National Average of Hospitals submitting data:

SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average



83

60

8

59

SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average

within 30 minutes of arrival Higher numbers are better		
Outpatients with chest pain or possible heart attack who got aspirin within 24 hours of arrival Higher numbers are better	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Outpatients having surgery who got an antibiotic at the right time - within one hour before surgery Higher numbers are better	Surgical Care Improvement Project	National Average of Hospitals submitting data:
Outpatients having surgery who got the right kind of antibiotic. Higher numbers are better	Surgical Care Improvement Project	National Average of Hospitals submitting data:
Median Time to Fibrinolysis	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Heart Attack Patients Given Aspirin at Arrival	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Attack Patients Given Fibrinolytic Medication Within 30 Minutes Of Arrival	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Attack Patients Given PCI Within 90 Minutes Of Arrival	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data scored equal to or higher than

SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average




SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average

Heart Attack Patients Given ACE Inhibitor or ARB for Left Ventricular Systolic Dysfunction (LVSD)	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Attack Patients Given Smoking Cessation Advice/Counseling	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Attack Patients Given Aspirin at Discharge	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Attack Patients Given Beta Blocker at Discharge	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Attack Patients Given a Prescription for a Statin at Discharge	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Failure Patients Given Discharge Instructions	Heart Failure	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Failure Patients Given an Evaluation of Left Ventricular Systolic (LVS) Function	Heart Failure	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Failure Patients Given ACE Inhibitor or ARB for Left Ventricular Systolic Dysfunction (LVSD)	Heart Failure	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Failure Patients Given Smoking Cessation Advice/Counseling	Heart Failure	Top 10% of Hospitals submitting data scored equal to or higher than

SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average




SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average

Advice/Counseling		
Pneumonia Patients Given Initial Antibiotic(s) within 6 Hours After Arrival	Pneumonia	Top 10% of Hospitals submitting data scored equal to or higher than
Pneumonia Patients Whose Initial Emergency Room Blood Culture Was Performed Prior To The Administration Of The First Hospital Dose Of Antibiotics	Pneumonia	Top 10% of Hospitals submitting data scored equal to or higher than
Pneumonia Patients Given the Most Appropriate Initial Antibiotic(s)	Pneumonia	Top 10% of Hospitals submitting data scored equal to or higher than
Pneumonia Patients Given Smoking Cessation Advice/Counseling	Pneumonia	Top 10% of Hospitals submitting data scored equal to or higher than
Pneumonia Patients Assessed and Given Pneumococcal Vaccination	Pneumonia	Top 10% of Hospitals submitting data scored equal to or higher than
Pneumonia Patients Assessed and Given Influenza Vaccination	Pneumonia	Top 10% of Hospitals submitting data scored equal to or higher than
Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent	Surgical Care Improvement Project	Top 10% of Hospitals submitting data scored equal to or higher than

SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average



SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average

infection

Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	Surgical Care Improvement Project	Top 10% of Hospitals submitting data scored equal to or higher than
Patients who got treatment at the right time (within 24 hours before or after their surgery) to help prevent blood clots after certain types of surgery	Surgical Care Improvement Project	Top 10% of Hospitals submitting data scored equal to or higher than
Surgery patients who were taking heart drugs called beta blockers before coming to the hospital, who were kept on the beta blockers during the period just before and after their surgery	Surgical Care Improvement Project	Top 10% of Hospitals submitting data scored equal to or higher than
Surgery patients who were given the right kind of antibiotic to help prevent infection	Surgical Care Improvement Project	Top 10% of Hospitals submitting data scored equal to or higher than
Heart surgery patients whose blood sugar (blood glucose) is kept under good control in the days right after surgery	Surgical Care Improvement Project	Top 10% of Hospitals submitting data scored equal to or higher than
Surgery patients needing hair removed from the surgical area	Surgical Care Improvement Project	Top 10% of Hospitals submitting data scored equal to or higher than

SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average

100

100

100

100

99

100

SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average


before surgery, who had hair removed using a safer method (electric clippers or hair removal cream – not a razor)		
Patients having surgery who were actively warmed in the operating room or whose body temperature was near normal by the end of surgery.	Surgical Care Improvement Project	Top 10% of Hospitals submitting data scored equal to or higher than
Surgery patients whose doctors ordered treatments to prevent blood clots after certain types of surgeries	Surgical Care Improvement Project	Top 10% of Hospitals submitting data scored equal to or higher than
Average number of minutes before outpatients with chest pain or possible heart attack who needed specialized care were transferred to another hospital A lower number of minutes is better	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data scored equal to or higher than
Average number of minutes before outpatients with chest pain or possible heart attack got an ECG A lower number of minutes is better	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data scored equal to or higher than
Outpatients with chest pain or	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data

SHG_POCCareMeasures_Nat


Based on Hospital Process of Care Measures – National Average



100



100



3

100

SHG_POCCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average

possible heart attack who got drugs to break up blood clots within 30 minutes of arrival
Higher numbers are better

scored equal to or higher than

Outpatients with chest pain or possible heart attack who got aspirin within 24 hours of arrival
Higher numbers are better

Heart Attack or Chest Pain

Top 10% of Hospitals submitting data scored equal to or higher than

Outpatients having surgery who got an antibiotic at the right time - within one hour before surgery
Higher numbers are better

Surgical Care Improvement Project

Top 10% of Hospitals submitting data scored equal to or higher than

Outpatients having surgery who got the right kind of antibiotic. Higher numbers are better

Surgical Care Improvement Project

Top 10% of Hospitals submitting data scored equal to or higher than

Median Time to Fibrinolysis

Heart Attack or Chest Pain

Top 10% of Hospitals submitting data scored equal to or higher than

SHG_POCareMeasures_Nat

Based on Hospital Process of Care Measures – National Average

100

100

100

18